

AOA CONCUSSION POLICY

Revised: February 9, 2020 by AOA Board of Directors

A concussion is a brain injury. All concussions should be regarded as potentially serious. Most people with concussions recover completely with correct management. However, incorrect management of a concussion can lead to further injury.

Concussions should be managed according to current **Canadian Harmonized Sport Concussion protocol** guidelines.

Concussions are to be diagnosed and managed by health care professionals working within their scope of practice and expertise.

Concussion symptoms must be completely resolved and medical clearance must be received before resuming any training. A progressive exercise program that re-introduces an individual to their full potential is recommended following concussion recovery.

What is a Concussion

Concussion is an injury to the brain resulting in a disturbance of brain function involving thinking and behavior.

What Causes a Concussion?

Concussion can be caused by a direct blow to the head or an impact to the body causing rapid movement of the head and subsequent movement of the brain within the skull.

Onset of Symptoms

Symptoms of concussion typically appear immediately but may evolve within the first 24-48 hours. For symptoms refer to: [Concussion Recognition Tool 5](#)

Who is at Risk?

Anyone.

- Children and adolescents (18 years and under) are more susceptible to brain injury, take longer to recover, and are susceptible to rare, dangerous brain complications, which may include death.
- Participants with previous concussion are at increased risk of further concussions.
- Awareness of the signs and symptoms of concussion and knowledge of how to properly manage a concussion is critical to recovery.



Policy

AOA is committed to maintaining the health of its participants and believes that a participant's health is more important than participating in sport events or activities. AOA recognizes the importance of increased awareness of concussions and their long-term effects and therefore enacts this policy as a tool to help manage concussed and possibly-concussed athletes.

This policy applies to all AOA organized or sanctioned events.

Procedure:

During all orienteering events, including competitions, training, or practice sanctioned by the AOA, all participants (including athletes, coaches, officials, or other members) will use their best efforts to be aware and take notice of incidents that may cause a concussion, and apply the AOA concussion protocol listed below.

AOA CONCUSSION PROTOCOL

Adapted from: Parachute. (2017). Canadian Guideline on Concussion in Sport.

www.parachute.ca/concussion

Purpose

This protocol covers the recognition, medical diagnosis, and management of all participants who may sustain a suspected concussion during an Orienteering /sport activity. It aims to ensure that athletes with a suspected concussion receive timely and appropriate care and proper management to allow them to return back to their sport safely. This protocol may not address every possible clinical scenario that can occur during sport-related activities but includes critical elements based on the latest evidence and current expert consensus.

Who should use this protocol?

This protocol is intended for use by all individuals who interact with athletes inside and outside the context of school and non-school based organized sports activity, including athletes, parents, coaches, officials, teachers, trainers, and licensed healthcare professionals.

For a summary of the AOA Sport Concussion Pathway refer to the diagram at the end of this document.

1. Education

Despite recent increased attention focusing on concussion there is a continued need to improve concussion education and awareness. Optimizing the prevention and management of concussion depends highly on annual education of all sport stakeholders (athletes, parents, coaches, officials, teachers, trainers, licensed healthcare professionals) on current evidence-informed approaches that can prevent concussion and more serious forms of head injury and help identify and manage an athlete with a suspected concussion.

AOA will provide resources in the form of online links and recommendations on its website, where its affiliated clubs and participants (which include coaches, athletes, officials, or other members) can learn more about concussion in sport. AOA will encourage its members to educate themselves about concussion in sport and will include information about the AOA concussion policy at coaching and official's courses. AOA recommends to its affiliated clubs to either adopt the AOA concussion policy or develop and implement their own concussion policy.

2. Head Injury Recognition

- ▶ **Refer to:** [Concussion Recognition Tool 5](#)

Although the formal diagnosis of concussion should be made following a medical assessment, all sport stakeholders including athletes, parents, teachers, coaches, teachers, officials, and licensed healthcare professionals are responsible for the recognition and reporting of athletes who may demonstrate visual signs of a head injury or who report concussion-related symptoms. This is particularly important because many sport and recreation venues will not have access to on-site licensed healthcare professionals.

A concussion should be suspected:

- in any athlete who sustains a significant impact to the head, face, neck, or body and demonstrates *ANY* of the visual signs of a suspected concussion or reports *ANY* symptoms of a suspected concussion as detailed in the *Concussion Recognition Tool 5*.
- if a player reports *ANY* concussion symptoms to one of their peers, parents, teachers, or coaches or if anyone witnesses an athlete exhibiting any of the visual signs of concussion.

If an athlete demonstrate signs or symptoms of a more severe head or spine injury including convulsions, worsening headaches, vomiting or neck pain. If an athlete demonstrates any of the 'Red Flags' indicated by the *Concussion Recognition Tool 5*, a more severe head or spine injury should be suspected, and Emergency Medical Assessment should be pursued.

- ▶ **Who:** Athletes, parents, coaches, officials, teachers, trainers, and licensed healthcare professionals
- ▶ **How:** [Concussion Recognition Tool 5](#)

3. Onsite Medical Assessment

Depending on the suspected severity of the injury, an initial assessment may be completed by emergency medical professionals or by an on-site licensed healthcare professional where available. In cases where an athlete loses consciousness or it is suspected an athlete might have a more severe head or spine injury, Emergency Medical Assessment by emergency medical professionals should take place (see 3a below). If a more severe injury is not suspected, the athlete should undergo Sideline Medical Assessment or Medical Assessment, depending on if there is a licensed healthcare professional present (see 3b below).

3a. Emergency Medical Assessment

If an athlete is suspected of sustaining a more severe head or spine injury during a game or practice, an ambulance should be called immediately to transfer the patient to the nearest emergency department for further Medical Assessment.

Coaches, parents, teachers, trainers and officials should not make any effort to remove equipment or move the athlete until an ambulance has arrived and the athlete should not be left alone until the ambulance arrives. After the emergency medical services staff has completed the Emergency Medical Assessment, the athlete should be transferred to the nearest hospital for Medical Assessment. In the case of youth (under 18 years of age), the athlete's parents should be contacted immediately to inform them of the athlete's injury. For athletes over 18 years of age, their emergency contact person should be contacted if one has been provided

- ▶ **Who:** Emergency medical professionals

3b. Sideline Medical Assessment

If an athlete is suspected of sustaining a concussion and there is no concern for a more serious head or spine injury, the player should be immediately removed from the field of play.

Scenario 1: If a licensed healthcare professional is present

The athlete should be taken to a quiet area and undergo Sideline Medical Assessment using the Sport Concussion Assessment Tool 5 (SCAT5) or the Child SCAT5. The SCAT5 and Child SCAT5 are clinical tools that should only be used by a licensed healthcare professional that has experience using these tools. It is important to note that the results of SCAT5 and Child SCAT5 testing can be normal in the setting of acute concussion. As such, these tools can be used by licensed healthcare professionals to document initial neurological status but should not be used

to make sideline return-to-sport decisions in youth athletes. **Any youth athlete who is suspected of having sustained a concussion must not return to the game or practice and must be referred for Medical Assessment.**

If a youth athlete is removed from play following a significant impact and has undergone assessment by a licensed healthcare professional, but there are NO visual signs of a concussion and the athlete reports NO concussion symptoms then the athlete can be returned to play but should be monitored for delayed symptoms.

In the case of national team-affiliated athletes (age 18 years and older), an experienced certified athletic therapist, physiotherapist or medical doctor providing medical coverage for the sporting event may make the determination that a concussion has not occurred based on the results of the Sideline Medical Assessment. In these cases, the athlete may be returned to the practice or **game without a *Medical Clearance Letter*** but this should be clearly communicated to the coaching staff. Players that have been cleared to return to games or practices should be monitored for delayed symptoms. If the athlete develops any delayed symptoms the athlete should be removed from play and undergo medical assessment by a medical doctor or nurse practitioner.

Scenario 2: If there is no licensed healthcare professional present

The athlete should be referred immediately for medical assessment by a medical doctor or nurse practitioner, and the athlete must not return to play until receiving medical clearance.

- ▶ **Who:** Athletic therapists, physiotherapists, medical doctor
- ▶ **How:** [*Sport Concussion Assessment Tool 5 \(SCAT5\)*](#), [*Child Sport Concussion Assessment Tool 5 \(Child SCAT5\)*](#)

4. Medical Assessment

In order to provide comprehensive evaluation of athletes with a suspected concussion, the medical assessment must rule out more serious forms of traumatic brain and spine injuries, must rule out medical and neurological conditions that can present with concussion-like symptoms, and must make the diagnosis of concussion based on findings of the clinical history and physical examination and the evidence-based use of adjunctive tests as indicated (i.e CT scan). In addition to nurse practitioners, medical doctors¹ that are qualified to evaluate patients with a suspected concussion include: pediatricians; family medicine, sports medicine,

¹ Medical doctors and nurse practitioners are the only healthcare professionals in Canada with licensed training and expertise to meet these needs; therefore all athletes with a suspected concussion should undergo evaluation by one of these professionals.

emergency department, internal medicine, and rehabilitation (physiatrists) physicians; neurologists; and neurosurgeons.

In geographic regions of Canada with limited access to medical doctors (i.e. rural or northern communities), a licensed healthcare professional (i.e. nurse) with pre-arranged access to a medical doctor or nurse practitioner can facilitate this role. The medical assessment is responsible for determining whether the athlete has been diagnosed with a concussion or not. Athletes with a diagnosed concussion should be provided with a *Medical Assessment Letter* indicating a concussion has been diagnosed.

Athletes that are determined to have not sustained a concussion must be provided with a *Medical Assessment Letter* indicating a concussion has not been diagnosed and the athlete can return to school, work and sports activities without restriction.

- ▶ **Who:** Medical doctor, nurse practitioner, nurse
- ▶ **How:** *Medical Assessment Letter*

5. Concussion Management

When an athlete has been diagnosed with a concussion, it is important that the athlete's parent/legal guardian is informed. All athletes diagnosed with a concussion must be provided with a standardized *Medical Assessment Letter* that notifies the athlete and their parents/legal guardians/spouse that they have been diagnosed with a concussion and may not return to any activities with a risk of concussion until medically cleared to do so by a medical doctor or nurse practitioner. Because the *Medical Assessment Letter* contains personal health information, it is the responsibility of the **athlete or their parent/legal guardian to provide** this documentation to the athlete's coaches, teachers, or employers. It is also important for the athlete to provide this information to sport organization officials that are responsible for injury reporting and concussion surveillance where applicable.

Athletes diagnosed with a concussion should be provided with education about the signs and symptoms of concussion, strategies about how to manage their symptoms, the risks of returning to sport without medical clearance and recommendations regarding a gradual return to school and sport activities. Athletes diagnosed with a concussion are to be managed according to their *Return-to-School and Sport-Specific Return-to-Sport Strategy* under the supervision of a medical doctor or nurse practitioner. When available, athletes should be encouraged to work with the team athletic therapist or physiotherapist to optimize progression through their *Sport-Specific Return-to-Sport Strategy*. **Once the athlete has completed their *Return-to-School and Sport-Specific Return-to-Sport Strategy* and are deemed to be clinically recovered from their concussion, the medical doctor or nurse practitioner can consider the athlete for a return to full sports activities and issue a *Medical Clearance Letter*.**

Return-to-School Strategy

The following is an outline of the *Return-to-School Strategy* that should be used to help student-athletes, parents, and teachers to collaborate in allowing the athlete to make a gradual return to school activities. Depending on the severity and type of the symptoms present student-athletes will progress through the following stages at different rates. If the student-athlete experiences new symptoms or worsening symptoms at any stage, they should go back to the previous stage. Athletes should also be encouraged to ask their school if they have a school-specific Return-to-Learn Program in place to help student-athletes make a gradual return to school.

Stage	Aim	Activity	Goal of each step
1	Daily activities at home that do not give the student-athlete symptoms	Typical activities during the day as long as they do not increase symptoms (i.e. reading, texting, screen time). Start at 5-15 minutes at a time and gradually build up.	Gradual return to typical activities
2	School activities	Homework, reading or other cognitive activities outside of the classroom.	Increase tolerance to cognitive work
3	Return to school part-time	Gradual introduction of schoolwork. May need to start with a partial school day or with increased breaks during the day.	Increase academic activities
4	Return to school full-time	Gradually progress	Return to full academic activities and catch up on missed school work

McCroory et al. (2017). Consensus statement on concussion in sport – the 5th international conference on concussion in sport held in Berlin, October 2016. *British Journal of Sports Medicine*, 51(11), 838-847.

AOA Return-to-Orienteering Strategy

The following is an outline of the Return-to-Sport Strategy that should be used to help athletes, coaches, trainers, and medical professionals to partner in allowing the athlete to make a gradual return to sport activities. An initial period of 24-48 hours of rest is recommended before starting the - ***AOA Return-to-Orienteering Strategy***. The athlete should spend a minimum duration of 24 hours without symptom increases at each stage before progressing to the next one. If the athlete experiences new symptoms or worsening symptoms at any stage, they should go back to the previous stage. It is important that youth and adult student-athletes return to full-time school activities before progressing to stage 5 and 6 of the ***AOA Return-to-Orienteering Strategy***. It is also important that all athletes provide their coach with a *Medical Clearance Letter* prior to returning to full contact sport activities.

AOA Return-to-Orienteering Strategy

Stage	Aim	Activity	Goal of each step
1	Symptom-limiting activity	Daily activities that do not provoke symptoms	Gradual re-introduction of work/school activities
2	Light aerobic activity	Walking or stationary cycling at slow to medium pace. No resistance training - <i>Light intensity jogging or stationary cycling for 15-20 minutes at sub-symptom threshold intensity</i> - <i>No map reading/navigational activities.</i>	Increase heart rate
3	Sport-specific exercise (map reading) at slow pace	Running drills. No head impact activities - <i>Moderate intensity jogging for 30-60 minutes at sub-symptom threshold intensity</i> - <i>Moderate difficulty of map reading/navigational activities with walking only</i>	Add movement
4	Full range of training with speed moderation of map reading activities	Harder training drills. May start progressive resistance training - <i>Participation in high intensity running and drills</i> - <i>Participation in resistance training work-outs</i> - <i>Moderate difficulty of map reading/navigational activities with slow pace running</i>	Exercise, coordination and increased thinking
5	Full range of practice	Following medical clearance - <i>Participation in full practice without activity restriction</i>	Restore confidence and assess functional skills by coaching staff
6	Return to sport	Normal competition participation	

McCrory et al. (2017). Consensus statement on concussion in sport – the 5th international conference on concussion in sport held in Berlin, October 2016. *British Journal of Sports Medicine*, 51(11), 838-847.

6. Multidisciplinary Concussion Care

Most athletes who sustain a concussion while participating in sport will make a complete recovery and be able to return to full school and sport activities within 1-4 weeks of injury. However, approximately 15-30% of individuals will experience symptoms that persist beyond this time frame. If available, individuals who experience persistent post-concussion symptoms (>4 weeks for youth athletes, >2 weeks for adult athletes) may benefit from referral to a medically supervised multidisciplinary concussion clinic that has access to professionals with licensed training in traumatic brain injury that may include experts in sport medicine, neuropsychology, physiotherapy, occupational therapy, neurology, neurosurgery, and rehabilitation medicine.

Referral to a multidisciplinary clinic for assessment should be made on an individualized basis at the discretion of an athlete's medical doctor or nurse practitioner. If access to a

multidisciplinary concussion clinic is not available, a referral to a medical doctor with clinical training and experience in concussion (e.g. a sport medicine physician, neurologist, or rehabilitation medicine physician) should be considered for the purposes of developing an individualized treatment plan. Depending on the clinical presentation of the individual, this treatment plan may involve a variety of health care professionals with areas of expertise that address the specific needs of the athlete based on the assessment findings.

- ▶ **Who:** Multidisciplinary medical team, medical doctor with clinical training and experience in concussion (e.g. a sports medicine physician, neurologist, or rehabilitation medicine physician), licensed healthcare professionals.


7. Return to Sport

Athletes who have been determined to have not sustained a concussion and those that have been diagnosed with a concussion and have successfully completed their *Return-to-School and AOA-Specific Return-to-Sport Strategy* can be considered for return to full sports activities. The final decision to medically clear an athlete to return to full game activity should be based on the clinical judgment of the medical doctor or nurse practitioner taking into account the athlete's past medical history, clinical history, physical examination findings and the results of other tests and clinical consultations where indicated (i.e. neuropsychological testing, diagnostic imaging). Prior to returning to full contact practice and competition, each athlete that has been diagnosed with a concussion **must provide their coach with a standardized *Medical Clearance Letter*** that specifies that a medical doctor or nurse practitioner has personally evaluated the patient and has cleared the athlete to return to sports. In geographic regions of Canada with limited access to medical doctors (i.e. rural or northern communities), a licensed healthcare professional (such as a nurse) with pre-arranged access to a medical doctor or nurse practitioner can provide this documentation. A copy of the *Medical Clearance Letter* should also be submitted to sports organization officials that have injury reporting and surveillance programs where applicable.

Athletes who have been provided with a *Medical Clearance Letter* may return to full sport activities as tolerated. If the athlete experiences any new concussion-like symptoms while returning to play, they should be instructed to stop playing immediately, notify their parents, coaches, trainer or teachers, and undergo follow-up *Medical Assessment*. In the event that the athlete sustains a new suspected concussion, the **AOA Concussion Protocol** should be followed as outlined here..

- ▶ **Who:** Medical doctor, nurse practitioner
- ▶ **Document:** *Medical Clearance Letter*

AOA Concussion Pathway



10* Persistent symptoms: lasting 2-3 weeks in children & youth or 2-4 weeks in adults